


CURRENT SCIENCE



Energy and environment: India

Large-scale renewable energy

Transportation: low-carbon CO₂ policy

Table 1. Shell weights (μg) of foraminifera species *Pulleniatina obliquiloculata*. CO_3^- ($\mu\text{mol}/\text{kg}$) calculated using the program developed by Takahashi; pressure-normalized carbonate ion concentration, $[\text{CO}_3^{*}]$ calculated from Broecker and Clark¹⁰, $\text{CO}_3^{*} = \text{CO}_3^- + 20(4 - h)$; where h is the depth (in km); ΔCO_3^- values calculated as $\text{CO}_3^{*} - \text{CO}_3^-$

Station	Depth (m)	Latitude	Longitude	Shell weight			
				<i>P. obliquiloculata</i>	CO_3^-	CO_3^{*}	ΔCO_3^-
SK199C/6	2250	8.13°N	73.56°E	33.17	77.59	112.59	35
SK199C/10	3305	7.36°S	67.17°E	29.11	83.61	97.51	13.9
SK199C/9	3320	4.86°S	67.09°E	32.05	83.70	97.29	13.6
SK199C/14	3368	15.27°S	66.01°E	35.1	83.97	96.61	12.64
SK199C/11	3373	9.17°S	65.95°E	31.63	83.99	96.54	12.54
SK199C/7	3944	5.51°N	69.34°E	26	87.25	88.37	1.12

bottom water ΔCO_3^- , from the Indian Ocean as given below.

$\Delta\text{CO}_3^- = [P. obliquiloculata \text{ shell weight } (\mu\text{g}) - 24.53]/0.2759$.

Using the above relation and shell weights of *P. obliquiloculata*, we calculated ΔCO_3^- interface. The gradient (ΔCO_3^- interface – ΔCO_3^- bottom water) was then calculated using, ΔCO_3^- bottom water values (Table 1).

Thus, we obtain a value of $+4.2 \mu\text{M}$ at 3900 m water depth. de Villiers¹² determined the ΔCO_3^- interface to be within $\pm 10 \mu\text{M}$ of bottom water. Shell weight measurements from both *Globigerinoides sacculifer* and *Globigerinoides ruber* suggest that there exists no significant gradient in ΔCO_3^- across the sediment–bottom water interface below a depth of 1000 m in the Indian Ocean. Our value of $+4.2 \mu\text{M}$ falls within the range of $\pm 10 \mu\text{M}$, suggesting that dissolution of carbonate around 3900 m water depth is due to undersaturation in CO_3^- concentration of bottom waters.

Calcium carbonate dissolution starts around 2250 m and intensifies at 3900 m in the tropical Indian Ocean⁸. We needed to study whether this intense dissolution occurring at 3900 m is a result of undersaturation of overlying bottom waters with respect to carbonate ion. This is important in order to know the calcite satu-

ration depth in the western tropical Indian Ocean, which is necessary to understand the oceanic response to the currently increasing CO_2 levels. We used the relationship between *P. obliquiloculata* shell weight and bottom water ΔCO_3^- from the core top data of Broecker and Clark¹¹ and data from the present study. We calculated ΔCO_3^- interface – ΔCO_3^- bottom water to be $+4.2 \mu\text{M}$, which is within $\pm 10 \mu\text{M}$ of bottom water, the value derived by de Villiers¹². This suggests that there is no gradient in ΔCO_3^- across the sediment–bottom water interface at a depth of 3900 m. Therefore, we conclude that the intense dissolution along this transect in the western tropical Indian Ocean is due to bottom water CO_3^- undersaturation.

- Berger, W. H., *Geol. Soc. Am. Bull.*, 1974, **18**, 1385–1402.
- Riley, J. P. and Chester, R., *Introduction to Marine Chemistry*, Academic Press, London, 1971, p. 465.
- Broecker, W. S. and Peng, T.-H., *Global Biogeochem. Cycles*, 1987, **1**, 15–29.
- Naidu, P. D. and Malmgren, B. A., *Mar. Geol.*, 1999, **161**, 49–62.
- Berger, W. H., *J. Foraminiferal Res.*, 1973, **3**, 187–195.
- Emerson, S. and Bender, M., *J. Mar. Res.*, 1981, **39**, 139–162.

- Archer, D. and Maier-Reimer, E., *Nature*, 1994, **367**, 260–263.
- Naik, S. S. and Naidu, P. D., *Geochem. Geophys. Geosyst.*, 2007, **8**, Q08009.
- Lohmann, G. P., *Paleoceanography*, 1995, **10**, 445–447.
- Broecker, W. S. and Clark, E., *Paleoceanography*, 1999, **14**, 596–604.
- Broecker, W. S. and Clark, E., *Paleoceanography*, 2001, **16**, 531–534.
- de Villiers, S., *Deep-Sea Res. I*, 2005, **52**, 671–680.

ACKNOWLEDGEMENTS. We thank Dr S. Chaturvedi for sharing SK samples. This is National Institute of Oceanography contribution no. 4366.

Received 5 November 2007; revised accepted 2 April 2008

S. S. NAIK¹
P. DIVAKAR NAIDU^{2,*}

¹National Centre for Antarctic and Ocean Research, Goa 403 804 India

²National Institute of Oceanography, Goa 403 004, India

*For correspondence.
e-mail: divakar@nio.org

Corallodiscus Batalin (Gesneriaceae): A new generic record for Eastern Ghats, Orissa

The floristics of southern Orissa, often considered incomplete, was sporadically approached by Mooney, Gamble, Haines and subsequent workers. Hence, it provides an ideal background for further exploration and discovery of taxonomic novelties. We collected an interesting specimen belonging to family Gesneriaceae

from Similipadar Hills (19°41'12.53"N, 83°4'28.97"E) in the Karlapat range, Kalahandi District, part of the Eastern Ghats, Orissa. Critical analysis and scrutiny of the floristic work done by several researchers^{1–8} helped us to identify this specimen as *Corallodiscus lanuginosus* (Wallich ex R.Br.) B. L. Burt. *C. lanuginosus* is a highly variable taxon and has been reported to grow in moist rock crevices at 600–3600 m amsl from northeast India to China. Recent discovery of this species from the Western Ghats⁸ and the Eastern Ghats (Prasad, RRL (B), col. no. 10217), extends its distribution to peninsular India. However, contrary to the



Figure 1. *Corallodiscus lanuginosus* (Wallich ex R.Br.) B. L. Burt (after Kamble *et al.*⁸).

claim of Kamble *et al.*⁸, Haines collected *Didissandra lanuginosa* var. *minuta* from Netarhat, Bihar, which was subsequently transferred to genus *Corallodiscus* Batalin by Naithani⁹. The genus *Corallodiscus* Batalin has 18 species described from the Himalayas to northwest China⁸. However, Zheng-Yi and Raven⁷ recognize 3–5 taxa in their study. The world check list for Gesneriaceae confirms five taxa under genus *Corallodiscus* Batalin, including *C. lanuginosus* var. *minuta* Naithani. Most of the species excluded were considered to be synonymous with *C. lanuginosus*, as the delimiting characters used for speciation are found to be continuous and justify a range. These variable characters are the indumentum (amount and type), lamina shape, size, margin and texture, and flower number and size. To facilitate easy identification, a description is provided here (Figure 1).

C. lanuginosus (Wallich ex R.Br.) B. L. Burt, *Gard. Chron. Ser. 3*, 122 (3180): 212. 1947; Hara, *Fl. E. Himal.* 298. 1966. *Didymocarpus lanuginosus* Wallich ex R.Br., *On Cyrtandreae* 118. 1839. *Didissandra lanuginosa* C.B. Clarke in A. DC. *Monogr. Phan.* 5: 66. 1833; C.B. Clarke in Hook. f. *Flora Br. India* 4:355.1884.

Herb, almost stemless. Leaves radical, rosette, exstipulate, sparsely hairy above, brown woolly on veins below; deep green above, velvet greenish-white below;

petiole 2.8–3 cm long, strong, sparsely long pubescent; lamina elliptic, ovate to obovate, obtuse, tapering towards base, crenate, 6.5–7 × 3.5–4 cm; palmately reticulate, veins prominent. Axillary long pedunculate lax cyme, 4–5 flowered, ca. 11 cm long, pink, hairs 1–2 mm, wavy. Flowers ca. 14 mm long, purplish-white, ebracteate, pedicels 7–9 mm long; calyx segments equal in size, sepals five, connate at base, imbricate, 2–3 × 1 mm, glabrescent, lanceolate, acute at apex; corolla tubular, purplish, ca. 12 mm long, bilipped, upper lip two-lobed, 1–2 mm long, margin reflexed, lower lip three-lobed, 3–4 mm long, woolly inside, stamens four, didynamous, epipetalous, longer ones 8–9 mm, shorter ones 4 mm, anthers dorsifixed, staminode one, 2 mm long; disc ring-like, orange; carpels 5 × 1 mm, glabrous, ovary 2 × 1 mm, unilocular, placenta-two, parietal, ovules on both sides of placenta; style 3 mm, slender, stigma bilobed; capsule glabrous, narrowly linear to elliptic, four-grooved, dehisces loculicidally; seeds minute, brown, tapering at both ends.

Flowering and fruiting: July–September.

Habitat: Grows on steep slopes in crevices of moist rocks at an altitude of 600 m.

Distribution: India (Simla, Dehradun, Sikkim, Maharashtra, Orissa, Bihar), Nepal, Bhutan, Thailand, Indo-China, Northwest China.

Specimens examined: Griffith-3835 (CAL), Smith-3345 (CAL), C.B. Clarke-7338 (CAL), Inayat-24824 (CAL), Smith and Cave-874, 822 (CAL).

Note: The present specimen completely differs in its fruit shape, arrangement of anthers and absence of bracts from those of the specimens examined. A critical taxonomical analysis is required for the taxon.

1. Saxena, H. O. and Brahmam, M., *Flora of Orissa*, 1994–96, p. 2.
2. Haines, H. H., *The Botany of Bihar and Orissa*, 1921–25, pp. 3–4.
3. Mooney, H. F., *Supplement to the Botany of Bihar and Orissa*, 1950.
4. Gamble, J. S., *Flora of Presidency of Madras*, 1921, p. 2.
5. Hooker, J. D., *Flora of British India*, 1885, p. 4.
6. Stainton, A., *Flowers of the Himalaya*, 2003.
7. Zheng-Yi, Wu and Raven, P. H. (eds), *Flora of China*, 1998, p. 18.
8. Kamble, Y. M., Shendage, S. M. and Yadav, S. R., *Rheedea*, 2006, **16**, 63–65.
9. Naithani, *Flowering Plants of India, Nepal & Bhutan*, 1990, p. 328.

ACKNOWLEDGEMENT. We thank Director, IMMT, Bhubaneswar for providing the necessary facilities and Dr Anton Weber, RBG, Edinburgh, for advice.

Received 13 November 2007; revised accepted 22 May 2008

NIRAD CHANDRA ROUT^{1,*}
NABIN KUMAR DHAL¹
PRASAD KUMAR DASH¹
ANIL KUMAR BISWAL²

¹*Institute of Minerals and Materials Technology (formerly Regional Research Laboratory), Bhubaneswar 751 013, India*

²*Department of Botany, North Orissa University, Baripada 757 003, India*

*For correspondence.
e-mail: ncrout@immt.res.in